Limits and Continuity (2/2)

Introduction to Engineering Mathematics

Prof. Joris Vankerschaver

Contents

- Infinite limits
- 2 Vertical and horizontal asymptotes
- 3 Limit laws
- 4 Continuity

Infinite limits

We say that f(x) has an **infinite limit** for $x \to a$ if the values of f(x) become **arbitrarily large** when $x \to a$.

We note

$$\lim_{x\to a} f(x) = \infty \quad \text{or} \quad \lim_{x\to a} f(x) = -\infty,$$

depending on the sign of f(x) near x = a.

- $\bullet \lim_{x \to 0} \frac{1}{x^2}$
- $\lim_{x\to 0+} \ln x$

Vertical asymptotes

The line x=a is a **vertical asymptote** of f(x) if at least one of the following is true:

$$\lim_{x\to a+} f(x) = \pm \infty \quad \text{or} \quad \lim_{x\to a-} f(x) = \pm \infty.$$

Example: find the vertical asymptotes of

- $y = \ln x$
- $y = \tan x$

Find all vertical asymptotes of the function

$$f(x) = \frac{|x-1|}{x^2 - 3x + 2}.$$

Hint: Look for zeros of the denominator that are not zeros of the numerator.

Limits at infinity (intuition)

What happens to f(x) when x becomes very large (positive/negative)?

Examples:

$$\lim_{x \to \pm \infty} \frac{x^2 - 1}{x^2 + 1}$$

 $\lim_{x \to +\infty} e^{-x}$

Limits at infinity (definition)

We write:

$$\lim_{x \to +\infty} f(x) = L$$

We mean:

- f(x) is defined on an interval $(a, +\infty)$
- f(x) gets as close as we want to L by taking x large enough.

$$\lim_{x\to -\infty} f(x) = M$$

- f(x) is defined on an interval $(-\infty, b)$
- f(x) gets as close as we want to M by taking x large enough (in the negative direction).

Horizontal asymptotes

The line y = L is a **horizontal asymptote** of f(x) if at least one of the following is true:

$$\lim_{x\to +\infty} f(x) = L \quad \text{or} \lim_{x\to -\infty} f(x) = L.$$

Examples: Find the horizontal asymptotes of

- $y = \frac{x^2 1}{x^2 + 1}$ $y = e^{-x}$
- $y = \frac{e^x e^{-x}}{e^x + e^{-x}}$

Limit laws

- $\mathbf{1} \lim_{x \to \infty} x = \infty$
- 3 $\lim_{x\to\infty}(f(x)+g(x))=\lim_{x\to\infty}f(x)+\lim_{x\to\infty}g(x)$
- $4 \lim_{x \to \infty} (cf(x)) = c \lim_{x \to \infty} f(x)$
- 6 $\lim_{x\to\infty} \frac{f(x)}{g(x)} = \frac{\lim_{x\to\infty} f(x)}{\lim_{x\to\infty} g(x)}$, if the denominator is not zero

Limit laws (continued)

- 7 $\lim_{x\to\infty} f(x)^{m/n} = \left(\lim_{x\to\infty} f(x)\right)^{m/n} = L^{m/n}$, if
 - $L \ge 0$ for n even
 - $L \neq 0$ for m negative
- 8 If $f(x) \leq g(x)$, then $\lim_{x \to \infty} f(x) \leq \lim_{x \to \infty} g(x)$
- $\lim_{x\to\infty} 1/x^n = 0, \text{ if } n>0$
- - c is constant (not equal to $\pm \infty$)
 - f is continuous (see later)

$$\lim_{x \to +\infty} \frac{2x^5 + 1}{x^5 + x^3 + 1}$$

$$\lim_{x \to -\infty} \frac{x^7}{\sqrt{x^{14} + 1}}$$

$$\lim_{x\to +\infty} \left(\sqrt{x^2+1}-x\right)$$

 $\lim_{x\to 0-}e^{1/x}$

 $\lim_{x\to +\infty}\sin x$

$$\lim_{x \to \pi/2-} e^{\tan x}$$

Infinite limits at infinity

Summary of limit techniques

- **1** Try "just substituting" $\pm \infty$
- 2 Use common manipulations:
 - Simplify
 - Highest powers
 - Conjugate
 - .

Special cases (to memorize):

$$\lim_{h\to 0}\frac{\sin h}{h}=1\quad \text{and}\quad \lim_{h\to 0}\frac{\cos h-1}{h}=0.$$

Continuity

- Intuitively, continuity = "no jumps"
- Mathematically, f(x) is **continuous** at an interior point x=a of its domain if

$$\lim_{x\to a} f(x) = f(a).$$

Example:
$$f(x) = \frac{x^2 - x - 3}{x - 2}$$
 is not continuous at $x = 2$.

Define $f(x)=\frac{x^2-x-2}{x-2}$ when $x\neq 2$ and f(x)=1 otherwise (i.e. f(2)=1). Is f(x) continuous at x=2?

Left/right continuity

At
$$x = a$$
, $f(x)$ is ...

- Right continuous if $\lim_{x\to a+} f(x) = f(a)$
- Left continuous if $\lim_{x \to a-} f(x) = f(a)$.

Example: the Heaviside function is defined by H(x)=0 when x<0 and H(x)=1 when $x\geq 0$. Is H(x) right or left continuous at x=0?

Continuity on an interval

The function f(x) is continuous on an interval $\left[a,b\right]$ if ...

- f(x) is continuous at every $x \in (a,b)$
- f(x) is right-continuous in a
- f(x) is left-continuous in b.

Example: Show that $f(x) = 1 - \sqrt{1 - x^2}$ is continuous on [-1, 1].

Making new continuous functions out of old ones

If f(x),g(x) are continuous at x=a, and c is a constant, then the following are also continuous at x=a:

- *f* + *g*
- *f* − *g*
- cf
- fg
- f/g (if $g(a) \neq 0$)

Which functions are continuous?

Continuous on their domain:

- Polynomials
- Rational functions
- Trigonometric functions + inverse trigonometric functions
- Square roots, *n*th roots
- log, exp

In particular:

- Polynomials are continuous for all $x \in \mathbb{R}$
- Rational functions P(x)/Q(x) are continuous for all x so that $Q(x) \neq 0$.

Composition of functions

If f(x) is continuous and $\lim_{x\to a} g(x) = b$, then

$$\lim_{x\to a} f(g(x)) = f(b) = f\left(\lim_{x\to b} g(x)\right).$$

Example:
$$\lim_{x\to 2} \sin\left(\frac{2-x}{4-x^2}\pi\right)$$

Corollary

If g is continuous at x=a and f is continuous at y=g(a), then $f\circ g$ is continuous at x=a.

Examples: Where are the following functions continuous?

- $h(x) = \sin x^2$
- $\bullet \ h(x) = \ln(1 + \cos x)$