

Integration: Partial Fractions

Introduction to Engineering Mathematics

Prof. Joris Vankerschaver

What are partial fractions?

Every rational function can be written as a sum of “simple” partial fractions. For example

$$\frac{x+2}{x^3-x} = \frac{-2}{x} + \frac{3}{2(x-1)} + \frac{1}{2(x+1)}.$$

In this lecture, we will find a recipe for the coefficients and terms in the partial fraction expansion.

Why are partial fractions useful?

The advantage is that the partial fractions are *much* easier to integrate:

$$\begin{aligned}\int \frac{x+2}{x^3-x} dx &= -2 \int \frac{dx}{x} + \frac{3}{2} \int \frac{dx}{x-1} + \frac{1}{2} \int \frac{dx}{x+1} \\ &= -2 \ln|x| + \frac{3}{2} \ln|x-1| + \frac{1}{2} \ln|x+1| + C.\end{aligned}$$

How to find the partial fraction expansion

Goal

To integrate **any rational function**: determine

$$\int \frac{P(x)}{Q(x)} dx = ???$$

where $P(x)$ and $Q(x)$ are polynomials.

Step 1: Divide if necessary

If the degree of $P(x)$ is greater than or equal to the degree of $Q(x)$, do a polynomial division:

$$\int \frac{P(x)}{Q(x)} dx = \int S(x) dx + \int \frac{R(x)}{Q(x)} dx,$$

with $S(x)$ the quotient and $R(x)$ the remainder.

- Recall: $\deg R(x) < \deg Q(x)$.
- From now on, we will suppose that this division has already been done, so that $\deg P(x) < \deg Q(x)$.

Special case 1: Linear denominator

If $Q(x)$ is a linear polynomial, i.e. $Q(x) = Ax + B$, then our integral takes the form

$$\int \frac{P(x)}{Q(x)} dx = \int \frac{K}{Ax + B} dx$$
$$= \dots$$

Special case 2: Quadratic denominator

If $Q(x)$ is a quadratic polynomial, then several cases are possible. After completing the square, we can have one of the following forms:

If $\deg P(x) = 0$:

- $\int \frac{dx}{x^2 - a^2}$
- $\int \frac{dx}{x^2 + a^2}$

If $\deg P(x) = 1$:

- $\int \frac{x dx}{x^2 - a^2}$
- $\int \frac{x dx}{x^2 + a^2}$

Step 2: Find the roots of the denominator

Do a factorization of the denominator $Q(x)$ into factors with **real** coefficients.

You will find:

- Some linear factors $(x - \alpha)$, with α roots of $Q(x)$
- Some quadratic factors $(Ax^2 + Bx + C)$ that cannot be further reduced.

Step 2: Find the roots of the denominator

Do a factorization of the denominator $Q(x)$ into factors with **real** coefficients.

You will find:

- Some linear factors $(x - \alpha)$, with α roots of $Q(x)$
- Some quadratic factors $(Ax^2 + Bx + C)$ that cannot be further reduced.

Important: Do not split into complex factors. For example:

$$x^3 + x = x(x^2 + 1).$$

Stop here, don't factor into $x(x + i)(x - i)$.

Case 2.1: Distinct roots

Assume that $Q(x) = (x - \alpha_1) \cdots (x - \alpha_k)$, with all α_i **distinct** and **real**.

Then the partial fraction expansion becomes

$$\frac{P(x)}{Q(x)} = \frac{A_1}{x - \alpha_1} + \frac{A_2}{x - \alpha_2} + \cdots + \frac{A_k}{x - \alpha_k},$$

where the coefficients A_1, \dots, A_k can be determined by adding the terms together and comparing with the left-hand side.

Example

Find $\int \frac{x+2}{x^3-x} dx$

Case 2.2: Irreducible quadratic factors

- For each quadratic factor, put a *linear term* in the numerator of the partial fraction.
- Deal with the linear factors as before.

Example

$$\begin{aligned}\frac{x+2}{x^3+x} &= \frac{x+2}{x(x^2+1)} \\ &= \frac{A}{x} + \frac{Bx+C}{x^2+1} \\ &= \dots\end{aligned}$$

Therefore $\int \frac{x+2}{x^3+x} dx = \dots$

Case 2.3: Repeated linear factors

- If $Q(x)$ has a repeated factor $(x - \alpha)^p$, then add p terms to the partial fraction expansion:

$$\frac{P(x)}{Q(x)} = \frac{A_1}{x - \alpha} + \frac{A_2}{(x - \alpha)^2} + \cdots + \frac{A_p}{(x - \alpha)^p} + [\text{other PFE}]$$

- Deal with other linear and quadratic factors as before.

Example

Determine the partial fraction expansion of $\frac{1}{x^2(x-1)^3}$.

Example

Find $\int \frac{dx}{x^3 - 5x^2 + 8x - 4}$.

Summary

- 1 If $\deg P(x) \geq \deg Q(x)$, do polynomial division.
- 2 Factor the denominator $Q(x)$ and write partial fractions for each root:

- Distinct roots (roots with multiplicity 1):

$$\text{PF} = \frac{A}{x - \alpha}.$$

- Irreducible quadratic factors:

$$\text{PF} = \frac{Ax + B}{x^2 + \dots}$$

- Root with multiplicity p :

$$\frac{A_1}{x - \alpha} + \dots + \frac{A_p}{(x - \alpha)^p}.$$

- 3 Find the coefficients in the partial fraction expansion by solving a system of equations.
- 4 Integrate the partial fraction expansion.